

Industry Update





SPECIAL GUEST

Hon. Emma McBride

Member for Dobell
Shadow Assistant Minister for Mental Health
Shadow Assistant Minister for Carers
Pharmacist






State of Play



12.36m COVID-19 vaccine primary doses not yet given

(Population 5 and over)



Booster & 5-11 yrs vaccinations coming soon

Source: DOH daily update and Guild calculations



2.3m children 5-11years to be vaccinated

5-11 year age population by jurisdiction (number)





1,740,000 vaccines in pharmacy

700,000
COVID-19 vaccines administered by community pharmacy reported in AIR

1,040,000
COVID-19 vaccines administered by community pharmacy reported in AIR

Number of pharmacies administering by state										
Vaccine	ACT	NSW	NT	QLD	SA	TAS	VIC	WA	TOTAL	
AstraZeneca	65	1113	19	731	272	88	693	435	3416	
Moderna	64	1074	17	703	262	87	683	416	3306	




Pfizer in Pharmacy

Last Week



Pfizer approved for use in pharmacies including booster program from 8 November for 18 years +

"It's a universal booster, meaning it's available for people who've had Pfizer, AstraZeneca or Moderna" Greg Hunt, Minister for Health 26 October 2021



Not too late to participate. Please refer to the 26 July email from Pharmacy Programs Administrator (PPA) or contact PPA: support@ppaonline.com.au



Ordering Pfizer

- If you responded to the simplified and accelerated EOI process from 26 July – nothing further required
- 1,156 pharmacies have ordered Pfizer
 - 927 pharmacies will receive stock by 12 Nov
 - 229 pharmacies will receive stock by 19 Nov
- Pharmacies are able to commence the administration of Pfizer doses as a primary course of vaccination - first or second dose - however are encouraged to focus efforts on the use of existing vaccine stock where possible





The Booster program

- Available from six months after completion of the primary course (second dose)
- AstraZeneca can also be used as a booster dose in the following situations:
 - received AstraZeneca for first two doses, if there are no contraindications or precautions for use
 - If a significant adverse reaction has occurred after a previous mRNA vaccine which contraindicates further doses of mRNA vaccine (e.g. anaphylaxis, myocarditis)
- ATAGI will make a recommendation on the potential use of Moderna as a booster down the track



After your Pfizer Vaccination – Patient Handout

- New information sheet now available
- Department Health Patient Handout available in multiple languages

COVID-19 vaccination resources including above available at: www.guild.org.au

Information for after your Pfizer (COMIRNATY®) COVID-19 Vaccine Today your community pharmacist has given you the Pfizer COVID-19 Vaccine. Congratulations on taking this vital step in protecting yourself, your family, and your community from COVID-19. Two doses of Pfizer COVID-19 Vaccine are needed for maximum protection, Pfizer COVID-19 Vaccine may also be used as a booster. If this is your first dose, the pharmacy staff can assist you with booking in for your second dose. If this is your second dose or booster, a digital vaccination certificate will be automatically generated for you by Services Australia. You can see your COVID-19 digital certificate in your Medicare account through the Express Plus Medicare app or myGov What to expect after vaccination You may experience some mild side effects following vaccination. Most side effects last no more than a couple of days and you will recover without any problems Common side effects after the Pfizer Less common side effects after the Pfizer COVID-19 Vaccine include: COVID-19 Vaccine include. · pain or swelling at the injection site · redness or itching at the injection site tiredness and fatigue enlarged lymph nodes (swelling or tenderness in the headache · feeling sick (nausea) or unwell muscle or joint pain and stiffness · fever and chills nain in the arm insomnia These side effects are usually mild, and start within a day after the vaccination and go away within one or two days. They can be managed by resting and using a pain medicine such as paracetamol or ibuprofen, or a cold compress if needed, Your Some people experience more significant flu-like symptoms and may require time away from normal activities. These symptom can occur after either dose but are more common after the second dose. If after a few days of the vaccine you have unexpected symptoms or you have an expected side effect listed above that has not gone away, you should seek medical advice from your doctor or healthcare professional. Be sure to tell them that you have recently received the Pfizer COVID-19 Vaccine Serious reactions such as allergic reactions are extremely rare and usually occur within 15 minutes of re which is why you are asked to wait in the pharmacy afterwards. If an allergic reaction occurs, your pharmacist is trained and equipped to deal with this. After you leave the pharmacy, call 000 if you think you are having an allergic reaction and experiencing severe symptoms such as difficulty breathing, wheezing, a fast heartbeat or collapsing Myocarditis and pericarditis are very rare side effects that may happen after receiving the Pfizer COVID-19 Vaccine. Myocarditis is inflammation of the heart and pericarditis is inflammation of the heart lining. In the rare cases it has occurred, it has been within two weeks of having the vaccine. It occurs more commonly after the second dose of the vaccine and in younger men under 30 years of age. Most cases of myocarditis or pericarditis linked to Pfizer COVID-19 Vaccine have been mild and recovery has been quick You should attend your nearest Emergency Department or call 000 if you have any of the following symptoms particularly within 1 to 5 days after vaccination · irregular or skipped heart beats or 'fluttering' shortness of breath or pain with breathing languages is available from www.health available from www.healtha The Pharmacy Australasian College Guild of Australia





Pharmacist Training

Pharmacist immunisers must undertake Commonwealth COVID-19 training

https://covid19vaccinationtraining.org.au

1. COVID-19: an introduction	4. Multi-dose vial training and delivery		
2. Handling and storage	5. Documentation and reporting		
3. Communication and purpose	6. Safety surveillance and reporting of adverse events following immunisation (AEFI		





Managing stock





Transferring of stock between pharmacies encouraged

Contact the VOC 1800 318 208 and provide details e.g. batch number, expiry date and number of doses

/

Taskforce can only accept 20 + vials of Astra Zeneca, and 10 + vials of Moderna per collection. Requests assessed for suitability considering shelf life, time in transit and clinic location

(

Once assessed you will receive a notification email confirming collection or disposal instructions

It's better to use a vial on a person (regardless of dose numbers) then have the vial expire in the fridge a few days later

Î

Please report wastage through CVAS



Vaccines

COVID Vaccines	5-11 years	12-17 years	18 years +	Booster 18 years +
AstraZeneca				Sometimes*
Moderna	2022			
Pfizer	2021			

^{*} AstraZeneca can be used as booster if received it for first two doses, if there are no contraindications or precautions for use, and if a significant adverse reaction has occurred after a previous mRNA vaccine which contraindicates further doses of mRNA vaccine (e.g. anaphylaxis, myocarditis)



Marketing Campaign Update

- Month to date there has been 52,943 digital vaccination bookings made directly onsite
- Should reach 2 million vaccinations administered in community pharmacy by end of November
- Digital ads have reached 783,937 people over the last week, for 16,299,632 impressions since the campaign commenced
- Month to date our advertising has been viewed 8,254,695 times by 2.3 million individuals
- This week ethnic radio goes live across
 29 community stations in all states
- Promotional packs will be sent out to member pharmacies next week




findapharmacy.com.au





Importance of checking AIR





COVID Point of Care Testing

- Support community pharmacy access to COVID-19 point of care tests approved by the TGA and included on the ARTG for:
 - provision of surveillance testing for pharmacy staff and asymptomatic patients
 - sale and supply of self-testing/home test kits to consumers
- Recognise PCR testing is the gold standard, however we support the use of other types of surveillance PoC testing in areas where COVID-19 is circulating in the community
- Support the provision of safe and quality PoC testing services
 consider the TGA guidance for implementing COVID-19
 RA PoC testing
- Acknowledge there may be pharmacies who do not wish to provide testing
- Support the sale and supply of self-testing kits to consumers from community pharmacy
 - o access to a health professional and staff who are all trained

Guild Position
Statement





COVID-19 Podcast

- Available online
- Natalie Willis, Chair of Policy & Regulation Sub Committee, National Councillor and WA Community Pharmacist
- Gerard Benedet, Queensland Branch Director







Latest Updates



Continued Dispensing Extensions

- Cth Continued Dispensing Emergency Measures extended until 30 June 2022
 - For Expanded list of S85 PBS medicines
 - Also requires S/T extension
- 13 Jan 2020 to 30 Jun 2021 360,000 people used CD
- Guild data shows range of health conditions
 - Diabetes, CVD, asthma/COPD, Parkinson's, epilepsy, glaucoma
- Need it to be permanent & to cover all people in need
 - CTG or S100 specialised medicines (e.g. HIV) not covered

Guild Digest

The Pharmacy
Guild of Australia

The **Guild Digest** is the pre-eminent publication examining the operation and financial performance of Australian community pharmacies. The Guild Digest is also used in negotiations with the Government for Community Pharmacy Agreements.

Be part of the 50th anniversary edition of the Guild Digest by clicking on the QR Code below to complete the survey







Workforce shortages

- We are aware there are significant challenges in recruiting/attracting high quality staff
- Guild is looking for evidence to assist in our advocacy work regarding visas and migrations to attract internationally trained pharmacists

If you have any evidence to help support our advocacy work, please contact your local Guild branch or send an email to nationalwr@guild.org.au



Indigenous Focus



Hesitancy Research – COVID PR Campaign

- Research shows Indigenous Australians reluctant to get a vaccine
- High percentage not booked in to be vaccinated and unlikely to do so
- More concerned about side effects and effectiveness
- Low trust in messaging/advertising
- Need to engage with key community leaders to educate



Source: CT Group, research conducted mid October





Resources for vaccination providers







Our Reconciliation Action Plan

Reconciliation Australia have recently approved our RAP which will now be formally launched this month.





End of Year Wrap Up



SPECIAL GUEST

TBC



8 December, 8pm AEDT



✓ Pre-registrations open now





Remember we are here for you!

Call 13 GUILD OR covid-19@guild.org.au